

Modern Physics Letters A
 © World Scientific Publishing Company

PARITY VIOLATION IN $d\vec{\gamma} \rightarrow np$ WITH PIONLESS EFFECTIVE THEORY

CHANG HO HYUN

*Department of Physics Education, Daegu University
 Gyeongsan 712-714, Republic of Korea,
 hch@daegu.ac.kr*

JAE WON SHIN

*Department of Physics, Sungkyunkwan University
 Suwon 440-746, Republic of Korea*

SHUNG-ICHI ANDO

*Theoretical Physics Group, School of Physics and Astronomy, The University of Manchester,
 Manchester M13 9PL, United Kingdom*

Received (Day Month Year)

Revised (Day Month Year)

We consider a pionless effective theory with dibaryon fields for the description of the weak process involving two nucleons. We construct leading order Lagrangians that contain nucleon-dibaryon weak coupling constants. We calculate the physical observable in the photodisintegration of the deuteron at threshold and obtain the result in terms of the nucleon-dibaryon weak coupling constants. Relation to existing calculations is discussed.

Keywords: Forces in hadronic systems and effective interactions.

PACS Nos.: 21.30.Fe

1. Introduction

It is quite recent that the effective field theory (EFT) has put its first step toward the realm of hadronic weak interaction.^{1,2} An experiment to measure parity-violation (PV) effects in polarized neutron capture by a proton at LANSCE and its upgrade at SNS have evoked interest in exploring the PV phenomena in few nucleon systems in the context of EFT.^{3,4,5} The results obtained thus far indicate that EFT is a working machine for the study of the PV processes; perturbative expansion converges well in both interactions and transition operators. Inspired by the precedent success of the EFT in PV, we try to formulate the PV EFT as simple as possible so that it will be feasible to explore various few nucleon PV phenomena with it. Pionless EFT with dibaryon fields has proven one of the most efficient EFTs applicable to few nucleon systems at low energies.^{6,7,8} PV EFT with dibaryon

2 *C. H. Hyun, J. W. Shin, S. Ando*

fields was already considered by Savage⁹, but only a part of more general pionless PV Lagrangian was accounted in the work. In this work, we construct pionless PV Lagrangian with dibaryon fields at leading order, and calculate the PV observable in $d\bar{\gamma} \rightarrow np$ at threshold. We discuss how the present result can be compared with previous theoretical calculations.

2. Theory

PV dibaryon-nucleon-nucleon (dNN) Lagrangian can be written as

$$\mathcal{L}_{\text{PV}}^{dNN} = \sum_{\Delta T} \mathcal{L}_{\text{PV}}^{\Delta T} \quad (1)$$

where ΔT is the isospin change at the dNN vertex. For the two nucleon system,

$$\mathcal{L}_{\text{PV}}^{dNN} = \mathcal{L}_{\text{PV}}^0 + \mathcal{L}_{\text{PV}}^1. \quad (2)$$

Because of the total angular momentum conservation, allowed transitions due to the PV interaction for the lowest angular momentum states are $^1S_0 \leftrightarrow ^3P_0$, and $^3S_1 \leftrightarrow ^1P_1$ due to $\mathcal{L}_{\text{PV}}^0$, and $^3S_1 \leftrightarrow ^3P_1$ due to $\mathcal{L}_{\text{PV}}^1$. Introducing dimensionless PV dNN coupling constants $h_{\text{dNN}}^{\Delta T}$, we have the PV dNN Lagrangians

$$\mathcal{L}_{\text{PV}}^0 = \frac{h_{\text{dNN}}^{0s}}{2\sqrt{2}\rho_d m_N^{5/2}} s_a^\dagger N^T \sigma_2 \sigma_i \tau_2 \tau_a \frac{i}{2} \left(\overleftrightarrow{\nabla} - \overleftarrow{\nabla} \right)_i N \quad (3)$$

$$+ \frac{h_{\text{dNN}}^{0t}}{2\sqrt{2}\rho_d m_N^{5/2}} t_i^\dagger N^T \sigma_2 \tau_2 \frac{i}{2} \left(\overleftrightarrow{\nabla} - \overleftarrow{\nabla} \right)_i N + \text{h.c.}, \quad (4)$$

for the isospin conserving part and

$$\mathcal{L}_{\text{PV}}^1 = i \frac{h_{\text{dNN}}^1}{2\sqrt{2}\rho_d m_N^{5/2}} \epsilon_{ijk} t_i^\dagger N^T \sigma_2 \sigma_j \tau_2 \tau_3 \frac{i}{2} \left(\overleftrightarrow{\nabla} - \overleftarrow{\nabla} \right)_k N + \text{h.c.} \quad (5)$$

for the isospin changing one. s_a and t_i are the two-nucleon dibaryon fields in 1S_0 and 3S_1 states, respectively. σ 's and τ 's in Eqs. (3,4,5) project the neutron-proton system into 3P_0 , 1P_1 and 3P_1 states, respectively.

3. Diagrams and Results

An observable is the PV photon polarization defined as

$$P_\gamma = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-} \quad (6)$$

where $\sigma_{+(-)}$ is the total cross section with the photon of helicity $+1$ (-1). With PV interactions, P_γ is given as¹⁰

$$P_\gamma = -2 \frac{\mathcal{M}_{E1}}{\mathcal{M}_{M1}}, \quad (7)$$




Fig. 1. Leading diagrams for $d\bar{\gamma} \rightarrow np$ process. Thick line represents the *dressed* dibaryon fields, thin line the nucleon and the wavy line the photon. Cross circle stands for PV dNN vertex, and the dNN vertices without any mark for the PC interaction.

where $\mathcal{M}_{M1}$ and $\mathcal{M}_{\tilde{E}1}$ are the parity-conserving (PC) M1 and parity-violating E1 amplitudes, respectively. PC M1 amplitude has been calculated very accurately in the pionless EFT with dibaryons^{6,7}, and we employ the result in this work.

Feynman diagrams for the process are depicted in Fig. 1. It has been shown in the literature that the PV polarization depends on isoscalar and isotensor components of the meson-exchange PV potential. In our framework, isospin conserving weak Lagrangians given by Eqs. (3,4) contribute to $\mathcal{L}_{\tilde{E}1}$. In the diagrams (a) and (b), we have the initial 3S_1 state, and $^1\tilde{P}_1$ state is admixed in the initial state by the weak Lagrangian $\mathcal{L}_{PV}^{0t}$, $|^1\tilde{P}_1\rangle \propto \mathcal{L}_{PV}^{0t}|^3S_1\rangle \propto h_{dNN}^{0t}$. Since the leading E1 transition operator conserves the total spin, we have the PV amplitude from the diagrams (a) and (b) as

$$\mathcal{M}_{a,b} = \langle ^1S_0 | \hat{E}_1 | ^1\tilde{P}_1 \rangle \propto h_{dNN}^{0t}, \quad (8)$$

where $\hat{E}_1$ is the E1 transition operator. For the diagram (c) we have a parity admixture in the final state, $|^3\tilde{P}_0\rangle \propto \mathcal{L}_{PV}^{0s}|^1S_0\rangle \propto h_{dNN}^{0s}$. We then have the PV transition amplitudes

$$\mathcal{M}_c = \langle ^3\tilde{P}_0 | \hat{E}_1 | ^3S_1 \rangle \propto h_{dNN}^{0s}, \quad (9)$$

We employ the convection current operator of the nucleon for $\hat{E}_1$, which is proportional to the nucleon momentum. Counting the order of on-shell momentum in each diagram, diagram (a) is suppressed to diagrams (b) and (c) by p^2 where p is the outgoing nucleon momentum. Consequently diagrams (b) and (c) give leading contributions, and we obtain the results in the $p \rightarrow 0$ limit as

$$i\mathcal{M}_b = \frac{a_0 \gamma h_{dNN}^{0t}}{6\rho_d m_N^{5/2}} \sqrt{\frac{\gamma\rho_d}{1-\gamma\rho_d}}, \quad i\mathcal{M}_c = \frac{a_0 \gamma h_{dNN}^{0s}}{6\rho_d m_N^{5/2}} \sqrt{\frac{\gamma\rho_d}{1-\gamma\rho_d}}. \quad (10)$$

a_0 is the scattering length for the 1S_0 scattering state, ρ_d is the effective range for the 3S_1 bound state, and $\gamma = \sqrt{m_N B}$, where B is the deuteron binding energy. With the PC M1 amplitude obtained from the same theory, we have

$$P_\gamma = -\frac{a_0 \gamma}{6\sqrt{2}\pi m_N \rho_d} \frac{h_{dNN}^{0s} + h_{dNN}^{0t}}{\kappa_1(1 - a_0 \gamma) - \frac{L_1}{2} a_0 \gamma^2}, \quad (11)$$

where $\kappa_1 = \mu_V/2 = 2.35$ and L_1 is a low energy constant fixed from the $np \rightarrow d\gamma$ total cross section at threshold.

4 C. H. Hyun, J. W. Shin, S. Ando

4. Discussion

Using the pionless EFT with dibaryon fields, we calculated the PV polarization in $d\bar{\gamma} \rightarrow np$ at threshold. Though the use of the EFT makes the calculation simple and easy, physical meaning of the result is not as transparent as the one obtained from the meson exchange picture of the weak NN potential, so called the DDH potential¹¹. In order to better figure out the physical meaning of the present result, and eventually make the EFT a useful method for PV phenomenology, it is necessary to have the relation between the “new” and “old” languages. There are several steps to establish the connection. First, we have to relate the weak meson-nucleon coupling constants in the DDH potential to the PV dNN vertices. Starting point may be the pionful EFT. Expanding the pionful PV potential to a certain order, and then integrating out the pion degree of freedom, we obtain the relation between pionful and pionless theories. Second, we have to be cautious in treating the EM operators for the external photons. Most of the calculation in the literature assumes Siegert’s theorem for the E1 operator. In Ref. 1, it was shown that the PV asymmetry in $\bar{n}p \rightarrow d\gamma$ with convection current (adopted in this work) overestimates the result with Siegert’s theorem by a factor of 2.9. It was shown in the same reference that the meson-exchange terms strongly cancel the convection term contribution, and thus make the net result similar to the one with Siegert’s theorem. Understanding the role of higher order operators may be crucial to put a bridge between pionful and pionless results. If the relation is well understood and thus the pionless EFT with dibaryon fields becomes an understandable language in the hadronic weak interaction, the method can be applied to a broader range of few-body PV phenomena.

Acknowledgments

Work of CHH was supported by the Daegu University Research Grant, 2008. Work of JWS was supported in part by the Korea Science and Engineering Foundation grant funded by the Korean Government (MOST) (No.M20608520001-07B0852-00110). SA is supported by STFC grant number PP/F000448/1.

References

1. C. H. Hyun, T.-S. Park and D.-P. Min, *Phys. Lett. B* **516**, 321 (2001).
2. S.-L. Zhu *et al.*, *Nucl. Phys. A* **748**, 435 (2005).
3. C. H. Hyun, S. Ando and B. Desplanques, *Phys. Lett. B* **651**, 257 (2007); *Eur. Phys. J. A* **32**, 513 (2007).
4. C.-P. Liu, *Phys. Rev. C* **75**, 065501 (2007).
5. B. Desplanques, C. H. Hyun, S. Ando and C.-P. Liu, *Phys. Rev. C* **77**, 064002 (2008).
6. S. Ando and C. H. Hyun, *Phys. Rev. C* **72**, 014008 (2005).
7. S. Ando, R. H. Cyburt, S. W. Hong and C. H. Hyun, *Phys. Rev. C* **74**, 025809 (2006).
8. S. Ando *et al.*, *Phys. Lett. B* **668**, 187 (2008).
9. M. J. Savage, *Nucl. Phys. A* **695**, 365 (2001).
10. B. R. Holstein, *Fizika B* **14**, 165 (2005).

11. B. Desplanques, J. F. Donoghue and B. R. Holstein, *Ann. Phys. (N.Y.)* **124**, 449 (1980).